



SteriTech Wound Protection Gel

A Liquid Plaster with Disinfecting Properties



Available sizes:
5mL Sachet; 250mL Push Disc
Dilution Factor:
Ready-to-use

Quick:

Dries to form a 'second skin' in 5-10 minutes depending on the type of wound.

Easy to Apply & Remove:

No distress to the wound bed & no training required to use it correctly.

Economical:

Saves on cost of plasters, bandages and dressings.

Residual Action:

Provides a Sterile dressing, and continues to disinfect even after skin has dried, meaning wounds stay contaminant free, giving long-lasting protection, and ultimate peace of mind.

Effective:

Keeps wounds moist, but contaminant free, resulting in accelerated healing, and decreased scar tissue formation.



Antimicrobial Efficacy:

SteriTech Concentrate has been tested effective against the following organisms by the SABS, hospitals, government approved & internationally accredited laboratories in South Africa, UK and the USA:

Bactericidal: *Pseudomonas aeruginosa*, *Staphylococcus aureus*, *Escherichia coli* & Coliforms, *Chlamydia trachomatis*, *Salmonella typhi* & *choleraesius*, *Enterobacter cloacae*, *Actinomyces naeslundii*, *Ureaplasma urealyticum*, *Streptococcus faecalis* & *mutans*, *Lactobacillus leichmanni* & *paracasei*, *Enterococcus hirae*, *Mycobacterium bovis* & *tuberculosis*, *Bacillus subtilis* & *cereus* and MRSA. **Virucidal:** Cytomegalovirus, HIV, Polio, Herpes, Influenza, Hepatitis, Rotavirus, Foot & Mouth Disease, Avian Influenza, SARS, Coronavirus, Norovirus, Parvo and Calicivirus. **Fungicidal** (including fungal spores): *Trichophyton mentagrophytes*, *Candida albicans* and *Aspergillus flavus* & *niger*.

Microbicidal: *Dermatophagoides pteronyssinus*

Active Ingredient:

SteriTech Wound Protection Gel contains SteriTech Concentrate (a Quaternary Ammonium Compound/TriButyl Tin Oxide Blend) at a level of 1.25%.

First Aid:

In case of eye contact:
Flush eyes thoroughly with water. Redness might occur, but should disappear within 30 minutes. Should it persist, consult your physician.

In case of ingestion:
Do not induce vomiting, simply drink a large glass of water and consult your physician.

Instructions for Use:

Apply directly from dispenser to wound site. Spread into as thin a layer as possible using a gloved finger, gauze or spatula. Allow 5-10 minutes to dry into a skin-like layer. **DO NOT COVER THE TREATED AREA WITH A BANDAGE OR PLASTER.** To remove, wet area & wipe lightly with a clean cloth, cotton wool or gauze. Reapply daily or after washing or bathing. For external use only.

Recommended Uses:

- Use to cover wounds in hard to reach places or where scarring is unwanted.
- Excellent on the following wound types: Bed Sores, Burns, Plaster Allergies, Fragile skin, Roasties, Cuts in hairline, Diabetic Ulcers
- Used as a liquid plaster/ second skin to keep wounds clean and bacteria free.

PPE Requirements:

Respiratory Protection: None
Ventilation: None
Eye Protection: None
Protective Gloves: None
Other Protective Clothing: None

Product Description:

Physical form: Clear Blue Paste
pH: 6 - 7
Rinsability: excellent
Odour: Mild chemical odour
Incompatibility: anionic detergents
natural soaps, strong oxidisers
Stability: Stable



Enquiries to:
1 Bohnvista, 25 Payne Street, Pinetown P.O. Box 1338, Kloof, 3640, South Africa Tel: +27 31 702 7656
Fax: +27 31 702 3659 Website: www.steritech.co.za Email: info@steritech.co.za

